Zoological Reports
An International Journal
OPEN ACCESS
Short Communication
Check for updates

Hydrothermal Systems and Their Role in the Chemical Origins of Life

Jelang Jelku D. Sangma[1,2,3]

[1]*Department of Food Science and Nutrition, University of Agricultural Sciences, GKVK, Bangalore, Karnataka, 560065, India*
[2]*AICRP-WIA, ICAR (CIWA), Bhubaneswar, College of Community Science, Central Agricultural University, Tura, Meghalaya - 794005, India*
[3]*Eudoxia Research University, 256 Chapman Road STE 105-4, New Castle, USA*

Corresponding author: **Jelang Jelku D. Sangma** | E-mail: **jelang.jelku3@gmail.com**

Citation: Jelang Jelku D. Sangma (2025). Hydrothermal Systems and Their Role in the Chemical Origins of Life. *Zoological Reports: An International Journal.* **DOI: https://doi.org/10.51470/ZR.2025.4.2.01**

Received 07 July 2025 | Revised 12 August 2025 | Accepted 10 September 2025 | Available Online October 17, 2025

Abstract

Hydrothermal systems have long been proposed as plausible environments for the chemical origins of life on early Earth. Characterized by dynamic physicochemical gradients, mineral-rich fluids, and sustained energy fluxes, these systems provide conditions conducive to prebiotic chemistry. Both deep-sea hydrothermal vents and terrestrial hot spring systems offer unique geochemical microenvironments that may have facilitated the synthesis, concentration, and organization of organic molecules into increasingly complex structures. Mineral surfaces, redox gradients, and naturally occurring proton gradients within hydrothermal settings are thought to have driven abiotic carbon fixation, polymerization reactions, and protometabolic pathways. Furthermore, compartmentalization within mineral pores may have acted as precursors to biological membranes, promoting molecular selection and catalytic enhancement. This review synthesizes current evidence regarding hydrothermal geochemistry, catalytic mineral surfaces, energy sources, and experimental simulations relevant to prebiotic chemistry. Competing models—including alkaline vent and surface hydrothermal field hypotheses—are critically evaluated. By integrating insights from geochemistry, molecular biology, and experimental origin-of-life research, this article highlights hydrothermal systems as dynamic reactors that potentially bridged the transition from geochemistry to biochemistry.

Keywords: *hydrothermal vents, origin of life, prebiotic chemistry, alkaline vents, mineral catalysis, protometabolism, chemical evolution, early Earth, redox gradients, abiogenesis.*

1. Introduction

The origin of life remains one of the most profound and enduring questions in science. Understanding how non-living chemical systems transitioned into self-sustaining biological entities requires identifying plausible early Earth environments capable of supporting complex prebiotic chemistry. Among the various hypotheses proposed—including primordial soup, ice matrix, and extraterrestrial delivery models—hydrothermal systems have emerged as leading candidates due to their rich chemical diversity and sustained energy availability. Early Earth, approximately 4.0–4.5 billion years ago, was geologically active, characterized by widespread volcanism, tectonic instability, and intense hydrothermal circulation. Interaction between seawater and hot basaltic crust generated hydrothermal vents on the ocean floor, while volcanic landmasses supported terrestrial hot springs and geysers [1]. These systems provided continuous fluxes of heat, reduced gases, metal ions, and mineral surfaces capable of catalyzing chemical reactions. Unlike static surface environments, hydrothermal systems generate strong physicochemical gradients—temperature, pH, redox potential, and ionic composition—that can drive non-equilibrium chemical processes. Modern biological systems universally rely on redox reactions and proton gradients to generate metabolic energy, suggesting that life may have harnessed similar geochemical gradients during its earliest stages [2]. Two principal hydrothermal models dominate current discussions: deep-sea alkaline hydrothermal vent systems and terrestrial hydrothermal fields. Each presents unique advantages for prebiotic chemistry, including catalytic mineral scaffolds, natural compartmentalization, and energy-rich conditions. Evaluating these systems provides insight into the chemical plausibility of life's emergence. This review examines the geological characteristics of hydrothermal systems, their chemical properties, experimental simulations, and their potential roles in facilitating key prebiotic processes such as carbon fixation, polymer formation, and compartmentalization.

2. Geological and Geochemical Characteristics of Hydrothermal Systems

Hydrothermal systems form when water interacts with heated rock, typically in tectonically active regions.

© ***Authors:*** *Published in* ***Zoological Reports: An International Journal*** *under the CC BY-NC-ND 4.0 license (http://creativecommons.org/licenses/by-nc-nd/4.0/). No commercial use or modifications permitted.*

In marine environments, seawater infiltrates oceanic crust, becomes superheated by underlying magma, and re-emerges enriched with dissolved metals and reduced compounds. These fluids can reach temperatures exceeding 350°C in black smoker systems, while alkaline vents typically operate at lower temperatures (40–90°C) [3]. Alkaline hydrothermal vents are particularly relevant to origin-of-life hypotheses. These systems produce hydrogen-rich, high-pH fluids through serpentinization reactions between water and ultramafic rocks. The resulting vent structures consist of porous mineral precipitates composed of iron-sulfur and carbonate minerals. These microporous matrices create natural compartments that separate vent fluids from surrounding ocean water, generating steep proton and redox gradients [4]. Terrestrial hydrothermal fields, such as hot springs and geysers, also exhibit fluctuating wet–dry cycles, temperature variation, and mineral-rich conditions. Periodic evaporation in these environments can concentrate organic molecules, facilitating polymerization reactions that might otherwise be inefficient in dilute oceanic conditions [5]. The geochemical diversity of hydrothermal systems provides reduced gases such as hydrogen, methane, ammonia, and hydrogen sulfide—key ingredients for prebiotic synthesis. Metal ions including iron, nickel, and molybdenum serve as catalytic centers for redox chemistry. These properties collectively make hydrothermal systems chemically favorable environments for prebiotic evolution.

3. Energy Sources and Redox Gradients

Life requires a continuous energy source to maintain chemical organization. Hydrothermal systems supply multiple energy forms, including thermal energy, redox potential differences, and natural electrochemical gradients. Redox gradients arise when reduced hydrothermal fluids encounter more oxidized ocean water. The resulting disequilibrium creates opportunities for electron transfer reactions, analogous to those driving modern cellular respiration. Hydrogen generated through serpentinization can serve as an electron donor, while carbon dioxide acts as an electron acceptor. Proton gradients across mineral barriers in alkaline vents resemble the proton motive force used by contemporary cells to synthesize ATP. Some researchers propose that early protometabolic systems exploited these naturally occurring gradients before evolving membrane-bound proton pumps [6]. The availability of continuous geochemical energy distinguishes hydrothermal systems from transient surface environments, making them attractive candidates for sustained chemical evolution.

4. Mineral Catalysis and Surface Chemistry

Mineral surfaces within hydrothermal systems provide catalytic platforms that enhance chemical reactions. Iron-sulfur minerals, such as greigite and pyrite, exhibit catalytic properties similar to active sites in modern enzymes. The structural resemblance between these minerals and the iron-sulfur clusters found in ancient metabolic enzymes supports the hypothesis of geochemical continuity. Surface adsorption concentrates organic molecules, increasing local reaction rates. Mineral pores create microenvironments that stabilize intermediates and promote polymer formation. Experimental studies demonstrate that amino acids and nucleotides can form under hydrothermal conditions and may polymerize when associated with mineral surfaces. Clay minerals and metal sulfides can facilitate condensation reactions necessary for forming peptides and nucleic acid chains [7]. These catalytic processes reduce the reliance on improbable spontaneous polymerization in dilute aqueous solutions.

5. Carbon Fixation and Protometabolism

One of the key challenges in origin-of-life research is explaining how simple carbon compounds transformed into complex biomolecules. Hydrothermal systems offer plausible pathways for abiotic carbon fixation through Fischer–Tropsch-type reactions and metal-catalyzed reduction of carbon dioxide. Experimental simulations show that iron and nickel catalysts under hydrothermal conditions can produce organic acids, hydrocarbons, and amino acid precursors. Some proposed protometabolic pathways resemble the reductive acetyl-CoA pathway, one of the most ancient carbon fixation routes in modern anaerobic microorganisms. The resemblance between hydrothermal chemistry and primitive metabolic networks suggests that life may have emerged from geochemically driven reaction cycles [8]. These protometabolic systems could have gradually incorporated organic catalysts, transitioning from mineral-based catalysis to enzyme-driven metabolism.

6. Compartmentalization and Proto-Cellular Structures

Compartmentalization is essential for concentrating molecules and sustaining reaction networks. In alkaline hydrothermal vents, microporous mineral structures provide natural compartments separated by thin inorganic walls. These compartments maintain pH and redox gradients, creating semi-isolated reaction chambers. Such mineral pores may have functioned as precursors to biological membranes [9], lipid-like amphiphilic molecules could have formed and assembled within these compartments, eventually replacing mineral boundaries with organic membranes. Terrestrial hydrothermal fields offer alternative compartmentalization mechanisms through wet–dry cycling. Repeated dehydration and rehydration promote vesicle formation and polymer stabilization. The integration of compartmentalization with energy gradients and catalytic surfaces strengthens the plausibility of hydrothermal origin scenarios.

Table 1: Important Features of Hydrothermal Systems Relevant to the Chemical Origins of Life

Hydrothermal Feature	Geochemical Characteristics	Prebiotic Chemical Role	Supporting Evidence	Implications for Early Life
Redox Gradients	Interaction of reduced vent fluids (H_2, CH_4, H_2S) with oxidized ocean water	Drives electron transfer reactions and abiotic carbon reduction	Laboratory simulations of CO_2 reduction using Fe–Ni catalysts	Foundation for protometabolic energy systems
Proton (pH) Gradients	Alkaline vent fluids (high pH) mixing with mildly acidic ocean water	Establishes natural electrochemical gradients similar to proton motive force	Microfluidic vent models sustaining proton differentials	Precursor to chemiosmotic energy conservation
Mineral Catalytic Surfaces	Iron-sulfur minerals, clays, metal sulfides	Catalyze organic synthesis and polymerization reactions	Experimental formation of amino acids and organic acids on mineral surfaces	Transition from geochemical to enzyme-based catalysis
Microporous Structures	Mineral chimney pores and carbonate compartments	Provide confinement, concentration, and selective diffusion	Microscopic analysis of vent chimneys showing compartmental networks	Early form of compartmentalization before lipid membranes
Thermal Gradients	Temperature variation from hot vent interiors to cooler surroundings	Facilitates molecular cycling, concentration, and reaction dynamics	Thermophoresis experiments concentrating nucleotides	Enhances molecular organization and chemical evolution
Continuous Energy Supply	Ongoing geothermal heat and serpentinization reactions	Sustains non-equilibrium chemical systems	Modern hydrothermal vent ecosystems and geochemical modeling	Supports sustained chemical evolution over geological timescales
Metal Ion Availability	Fe, Ni, Mo, Co dissolved in vent fluids	Act as catalytic cofactors in redox reactions	Structural similarity between Fe–S minerals and ancient enzyme clusters	Evolutionary continuity between mineral and biological catalysts

7. Experimental Simulations and Laboratory Evidence

Laboratory experiments replicating hydrothermal conditions provide empirical support for prebiotic synthesis. High-pressure reactors and microfluidic systems simulate vent environments, demonstrating formation of amino acids, lipids, and nucleotide precursors. Microcompartment models have shown that mineral membranes can sustain proton gradients for extended periods. Other experiments reveal that thermal gradients can drive molecular concentration through thermophoresis. While challenges remain—particularly regarding RNA stability at high temperatures—moderate-temperature alkaline vents offer conditions compatible with organic molecule persistence [10].

8. Competing Hypotheses and Critical Evaluation

Despite strong support for hydrothermal models, alternative hypotheses remain viable. Surface pond scenarios emphasize ultraviolet-driven synthesis and wet–dry cycles. Ice matrix models highlight protective freezing environments [11]. Hydrothermal systems must address concerns regarding molecular stability at elevated temperatures and dilution effects in oceanic settings. However, spatial heterogeneity within vents and existence of cooler microenvironments mitigate these concerns. Rather than being mutually exclusive, surface and hydrothermal models may represent complementary stages in chemical evolution.

9. Implications for Astrobiology

Hydrothermal systems are not unique to Earth. Evidence of subsurface oceans and hydrothermal activity on icy moons such as Europa and Enceladus suggests that similar processes may operate elsewhere in the solar system. The universality of water–rock interactions and redox gradients implies that hydrothermal environments could represent common habitats for life's emergence [12-13]. Studying terrestrial hydrothermal systems thus informs the search for extra-terrestrial life.

10. Conclusion

Hydrothermal systems represent one of the most compelling and scientifically coherent models for the chemical origins of life on early Earth. Their significance lies not in a single favorable factor, but in the convergence of multiple complementary processes within a single dynamic environment. These systems provided sustained energy flux through redox disequilibria, naturally occurring proton gradients, catalytic mineral surfaces, and structurally complex microporous compartments. Together, these features created localized conditions capable of driving chemical reactions far from thermodynamic equilibrium—an essential requirement for the emergence of organized, self-sustaining systems. The geochemical processes occurring within alkaline hydrothermal vents, particularly serpentinization-driven hydrogen production and iron–sulfur mineral formation, closely parallel core aspects of ancient metabolic pathways. The structural resemblance between mineral catalysts and modern enzyme cofactors suggests a plausible evolutionary continuity from inorganic geochemistry to biological catalysis. Furthermore, naturally generated electrochemical gradients across mineral barriers provide a mechanistic bridge to chemiosmotic energy conservation, a universal feature of life today. Terrestrial hydrothermal systems add another dimension by introducing wet–dry cycling and concentration mechanisms that may have facilitated polymerization and membrane assembly, marine and terrestrial hydrothermal models may represent complementary stages in chemical evolution, contributing distinct yet synergistic processes toward increasing molecular complexity and regarding the stability of informational polymers and the transition to fully autonomous cellular systems—ongoing experimental simulations, microfluidic vent models, and geochemical reconstructions continue to strengthen the plausibility of hydrothermal origin scenarios. Importantly, hydrothermal environments are not unique to Earth; similar water–rock interactions likely occur on icy moons and early planetary bodies, expanding the relevance of this model to astrobiology, hydrothermal systems provide an integrated framework linking planetary geodynamics, prebiotic chemistry, protometabolism, and early bioenergetics.

Acknowledgement

The author(s) gratefully acknowledge the support of the International Innovation Program for Post-Doctoral Fellow, Eudoxia Research University, USA and India, under registration ID ERU/IIP-PDF/REG/2024/293, for providing research support and academic resources that contributed to the completion of this work.

References

1. Holm, N. G. (1992). Why are hydrothermal systems proposed as plausible environments for the origin of life?. In *Marine Hydrothermal Systems and the Origin of Life: Report of SCOR Working Group 91* (pp. 5-14). Dordrecht: Springer Netherlands.
2. Martin, W., Baross, J., Kelley, D., & Russell, M. J. (2008). Hydrothermal vents and the origin of life. *Nature Reviews Microbiology*, *6*(11), 805-814.
3. Holm, N. G., & Andersson, E. (2005). Hydrothermal simulation experiments as a tool for studies of the origin of life on earth and other terrestrial planets: a review. *Astrobiology*, *5*(4), 444-460.
4. Chatterjee, S. (2016). A symbiotic view of the origin of life at hydrothermal impact crater-lakes. *Physical Chemistry Chemical Physics*, *18*(30), 20033-20046.
5. Macleod, G., McKeown, C., Hall, A. J., & Russell, M. J. (1994). Hydrothermal and oceanic pH conditions of possible relevance to the origin of life. *Origins of Life and Evolution of the Biosphere*, *24*(1), 19-41.
6. Martin, W., & Russell, M. J. (2007). On the origin of biochemistry at an alkaline hydrothermal vent. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *362*(1486), 1887-1926.
7. Oparin, A. I. (1969). Chemistry and the origin of life. *Royal Institute of Chemistry, Reviews*, *2*(1), 1-12.
8. Sojo, V., Herschy, B., Whicher, A., Camprubi, E., & Lane, N. (2016). The origin of life in alkaline hydrothermal vents. *Astrobiology*, *16*(2), 181-197.
9. Sojo, V., Herschy, B., Whicher, A., Camprubi, E., & Lane, N. (2016). The origin of life in alkaline hydrothermal vents. *Astrobiology*, *16*(2), 181-197.
10. Krishnamurthy, R., & Hud, N. V. (2020). Introduction: Chemical evolution and the origins of life. *Chemical Reviews*, *120*(11), 4613-4615.
11. Stüeken, Eva Elisabeth, R. E. Anderson, J. S. Bowman, W. J. Brazelton, J. Colangelo-Lillis, A. D. Goldman, S. M. Som, and J. A. Baross. "Did life originate from a global chemical reactor?." *Geobiology* 11, no. 2 (2013): 101-126.
12. Ruiz-Mirazo, K., Briones, C., & de la Escosura, A. (2014). Prebiotic systems chemistry: new perspectives for the origins of life. *Chemical reviews*, *114*(1), 285-366.
13. Herschy, Barry, Alexandra Whicher, Eloi Camprubi, Cameron Watson, Lewis Dartnell, John Ward, Julian RG Evans, and Nick Lane. "An origin-of-life reactor to simulate alkaline hydrothermal vents." *Journal of molecular evolution* 79, no. 5 (2014): 213-227.